

# Variational estimates for martingale transforms

Pavel Zorin-Kranich

University of Bonn

2020-06-09

Joint work with P. Friz.

# Rough paths

## Definition

A  $p$ -rough path,  $2 < p < 3$ , is a pair

$$X : [0, \infty) \rightarrow H, \quad \mathbb{X} : \Delta = \{(s, t) \mid 0 \leq s < t < \infty\} \rightarrow H \otimes H$$

such that  $X \in V_{\text{loc}}^p$ ,  $\mathbb{X} \in V_{\text{loc}}^{p/2}$ , and for  $s < t < u$

$$\mathbb{X}_{s,u} = \mathbb{X}_{s,t} + \mathbb{X}_{t,u} + (X_u - X_t) \otimes (X_t - X_s). \quad (\text{Chen's relation})$$

$p$ -variation:

$$V^p X = \sup_{l_{\max}, u_0 < \dots < u_{l_{\max}}} \left( \sum_{l=1}^{l_{\max}} |X_{u_l} - X_{u_{l-1}}|^p \right)^{1/p},$$

$$V^p \mathbb{X} = \sup_{l_{\max}, u_0 < \dots < u_{l_{\max}}} \left( \sum_{l=1}^{l_{\max}} |\mathbb{X}_{u_{l-1}, u_l}|^p \right)^{1/p}.$$

How to check the conditions  $X \in V^p$  and  $\mathbb{X} \in V^{p/2}$ ?

# Rough path lifts of martingales

Let  $M = (M_t)$  be a (Hilbert space valued) càdlàg martingale. Let

$$\mathbb{M}_{s,t} := \int_{(s,t]} (M_{u-} - M_s) \otimes dM_u.$$

Then, a.s., the pair  $(M, \mathbb{M})$  is a  $p$ -rough path for any  $p > 2$ .

- ▶ Chen's relation – from Itô integration
- ▶ Bound for  $V^p M$ : Lépingle 1976.
- ▶ Bounds for  $V^{p/2} \mathbb{M}$ :
  - ▶  $M$  Brownian motion: Lyons 1998
  - ▶  $M$  has continuous paths: Friz+Victoir 2006
  - ▶  $M$  dyadic: Do+Muscalu+Thiele 2010,
  - ▶  $M$  has càdlàg paths: Chevyrev+Friz 2017, Kovač+ZK 2018.

There exist rough path lifts of over processes, e.g. Lévy processes.

Q: what is the appropriate generality for these lifting results?

How to incorporate e.g. fractional Brownian motion?

## Joint rough path lifts

All martingales and processes are adapted, càdlàg, Hilbert space valued.

**Theorem (Friz+ZK 2020+)**

Let  $M = (M_t)$  be a càdlàg martingale and  $(X, \mathbb{X})$  a deterministic càdlàg  $p$ -rough path ( $2 < p < 3$ ). Then, a.s., the pair of processes

$$\begin{pmatrix} X \\ M \end{pmatrix}, \begin{pmatrix} \mathbb{X} & \int X_{u-} \otimes dM_u \\ \int M_{u-} \otimes dX_u & \int M_{u-} \otimes dM_u \end{pmatrix}$$

is a  $p$ -rough path.

Motivation:  $dY = a(Y) d\mathbf{X} + b(Y) dM$

New in this result:

- ▶ Variational estimates for Itô integrals  $\int X dM$ ,
- ▶ existence of and estimates for integrals  $\int M dX$ .

The proof also recovers existence of Itô integrals and estimates for  $\mathbb{M} = \int M dM$  from previous slide.

# Martingale transforms

Let  $(f_n)_{n \in \mathbb{N}}$  be a discrete time adapted process  
and  $(g_n)_{n \in \mathbb{N}}$  a discrete time martingale. Define *paraproduct*

$$\Pi_{s,t}(f, g) := \sum_{s < j \leq t} (f_{j-1} - f_s) dg_j, \quad dg_j = g_j - g_{j-1}.$$

Martingale in  $t$  variable, discrete version of area integral.

## Theorem (Main estimate)

Let  $1 \leq p \leq \infty$ ,  $0 < q_1 \leq \infty$ ,  $1 \leq q_0 < \infty$ . Define  $q$  by  $1/q = 1/q_0 + 1/q_1$  and suppose  $1/r < 1/2 + 1/p$ . Then, with  $\|g\|_{L^q} = (\mathbb{E}|g|^q)^{1/q}$ ,  $Sg = [g]^{1/2}$ ,

$$\left\| V^r \Pi(f, g) \right\|_{L^q} \lesssim \sup_{\tau} \left\| \left( \sum_k \left( \sup_{\tau_{k-1} < j \leq \tau_k} |f_{j-1} - f_{\tau_{k-1}}| \right)^p \right)^{1/p} \right\|_{L^{q_1}} \|Sg\|_{L^{q_0}}.$$

The supremum is taken over increasing sequences of stopping times  $\tau = (\tau_k)$ .

- ▶ If  $f$  is a martingale,  $p = 2$ ,  $1 \leq q_1 < \infty$ , then by BDG inequality the RHS is  $\lesssim \|Sf\|_{L^{q_0}} \|Sg\|_{L^{q_0}}$ . In this case, any  $r > 1$  works.
- ▶ For general  $f$ , RHS is  $\leq \|V^p f\|_{L^{q_0}} \|Sg\|_{L^{q_0}}$  and  $r = p/2$  works.

# Discrete approximation of adapted processes

## Definition

An *adapted partition*  $\pi = (\pi_j)_j$  is an increasing sequence of stopping times. Adapted partitions are ordered by a.s. inclusion of the sets  $\{\pi_j \mid j \in \mathbb{N}\}$ . The set of adapted partitions is directed, so  $\lim_{\pi}$  makes sense. For an adapted partition  $\pi$ , let

$$[t, \pi] := \max\{s \in \pi \mid s \leq t\}, \quad f_t^{(\pi)} := f_{[t, \pi]}.$$

## Lemma

If  $f \in L^q(V^p)$  for some  $q > 0$  and  $p > 1$ , then

$$\lim_{\pi} f^{(\pi)} = f \quad \text{in} \quad L^q(V^{\tilde{p}})$$

for any  $\tilde{p} \in (p, \infty) \cup \{\infty\}$ .

## Proof.

Given  $\epsilon > 0$ , consider the adapted partition

$$\pi_0 := 0, \quad \pi_{j+1}(\omega) := \inf\{t > \pi_j(\omega) \mid |f_t - f_{\pi_j(\omega)}|(\omega) > \epsilon\}.$$

□

# Stopping time reduction

$f$  adapted process,  $g$  martingale

Martingale transform:  $\Pi_{s,t}(f, g) = \sum_{s < j \leq t} (f_{j-1} - f_s) dg_j$

Square function:  $Sg = [g]^{1/2}$ , Hölder exponents:  $1/q = 1/q_0 + 1/q_1$ .

## Theorem (Main estimate)

Suppose  $1/r < 1/p + 1/2$ . Then

$$\|V^r \Pi\|_{L^q(\Omega)} \lesssim \|V^p f\|_{L^{q_1}(\Omega)} \|Sg\|_{L^{q_0}(\Omega)}.$$

The  $V^r$  norm is estimated as follows.

## Lemma

Let  $(\Pi_{s,t})_{s \leq t}$  be a càdlàg adapted sequence with  $\Pi_{t,t} = 0$  for all  $t$ .

Then, for every  $0 < \rho < r < \infty$  and  $q \in (0, \infty]$ , we have

$$\|V^r \Pi\|_{L^q} \lesssim \sup_{\tau} \left\| \left( \sum_{j=1}^{\infty} \left( \sup_{\tau_{j-1} \leq t < t' \leq \tau_j} |\Pi_{t,t'}| \right)^{\rho} \right)^{1/\rho} \right\|_{L^q}, \quad (1)$$

where the supremum is taken over all adapted partitions  $\tau$ .

# Lépingle's inequality

Above stopping time argument first used in the following result.

## Theorem (ZK 2019)

Let  $M$  be a martingale and  $w$  a positive random variable.

For  $1 < p < \infty$  and  $2 < r$ , we have

$$\|V^r M\|_{L^p(w)} \leq C_{p,r} A_p(w)^{\max(1, 1/(p-1))} \|M\|_{L^p(w)},$$

where the  $A_p$  characteristic is given by

$$A_p(w) := \sup_{\tau \text{ stopping time}} \|\mathbb{E}(w \mid \mathcal{F}_\tau) \mathbb{E}(w^{-1/(p-1)} \mid \mathcal{F}_\tau)^{p-1}\|_{L^\infty(w)}$$

Classical Lépingle's inequality is the case  $w \equiv 1$ ,  $A_p(w) = 1$ .

Weighted inequalities imply vector-valued inequalities.

For dealing with martingale transforms, we use vector-valued BDG inequalities that follow from weighted inequalities by Osękowski.



# Integration by parts

$(X, \mathbb{X})$  rough path,  $M$  martingale

So far we have estimated  $\int X dM$  and  $\int M dM$ .

Next, we want to construct and estimate  $\Pi(M, X) = \int M dX$ .

We do this by partial integration:

$$\Pi(M, X) := \delta M \delta X - \Pi(X, M) - \delta[X, M].$$

The bracket is given by

$$[X, M]_T = \sum_{u \leq T} \Delta X_u \Delta M_u, \quad \Delta M_u = M_u - M_{u-}.$$

Variation norm estimate for the bracket:

$$\begin{aligned} \|V^r[X, M]\|_{L^q} &\stackrel{\text{stopping}}{\lesssim} \left\| \left( \sum_{j=1}^{\infty} \left( \sup_{\tau_{j-1} < t < t' \leq \tau_j} |\delta[X, M]_{t,t'}| \right)^\rho \right)^{1/\rho} \right\|_{L^q} \\ &\stackrel{\text{vector BDG}}{\lesssim} \left\| \left( \sum_{j=1}^{\infty} \left( \sum_{\tau_{j-1} < u \leq \tau_j} |\Delta X_u \Delta M_u|^2 \right)^{\rho/2} \right)^{1/\rho} \right\|_{L^q} \\ &\stackrel{\text{Hölder}}{\leq} V^p X \cdot \left\| \left( \sum_{j=1}^{\infty} \sum_{\tau_{j-1} < u \leq \tau_j} |\Delta M_u|^2 \right)^{1/2} \right\|_{L^q} \end{aligned}$$